

TECHNION-PH-2009-08

UdeM-GPP-TH-09-178

EFI 09-15

arXiv:0905.1495 [hep-ph]

May 2009

DIAGNOSTIC FOR NEW PHYSICS IN $B \rightarrow \pi K$ DECAYS¹

Seungwon Baek

*Institute of Basic Science and Department of Physics**Korea University, Seoul 136-701, Korea*

Cheng-Wei Chiang

*Department of Physics and Center for Mathematics and Theoretical Physics**National Central University, Chungli, Taiwan 320, R.O.C. and**Institute of Physics, Academia Sinica, Taipei, Taiwan 115, R.O.C.*

Michael Gronau

Physics Department, Technion, Haifa 32000, Israel.

David London

*Physique des Particules, Université de Montréal**C. P. 6128, succ. centre-ville, Montréal, QC, Canada H3C 3J7*

Jonathan L. Rosner

*Enrico Fermi Institute and Department of Physics, University of Chicago**Chicago, IL 60637, U.S.A.*

A recent analysis of $B \rightarrow \pi K$ decays concludes that present data do not clearly indicate whether (i) the standard model (or $\Delta I = 0$ new physics) is sufficient, or (ii) $\Delta I = 1$ new physics is needed. We show that these two possibilities can be distinguished by whether a sum rule relating the CP asymmetries of the four $B \rightarrow \pi K$ decays is valid. If case (i) is favored, the sum rule holds, and one predicts $A_{CP}(\pi^0 K^0) = -0.15$, while in case (ii) fits to new physics involving large values of a color-suppressed tree amplitude entail $A_{CP}(\pi^0 K^0) = -0.03$. The current experimental average $A_{CP}(\pi^0 K^0) = -0.01 \pm 0.10$ must be measured a factor of at least three times more precisely in order to distinguish between the two cases.

PACS codes: 12.15.Hh, 12.15.Ji, 13.25.Hw, 14.40.Nd

¹To be submitted to Physics Letters B.

Several suggestions have been made in the past few years that data for $B \rightarrow \pi K$ decays cannot be simply explained within the Cabibbo-Kobayashi-Maskawa (CKM) framework. In most cases these claims suffered from uncertainties in QCD calculations of hadronic matrix elements using a heavy-quark expansion. The purpose of this note is to outline some of these claims, proposing a way for using improved data in order to make a more robust statement about new physics.

The four $B \rightarrow \pi K$ decay amplitudes are related by isospin [1],

$$A(B^0 \rightarrow \pi^- K^+) - \sqrt{2}A(B^+ \rightarrow \pi^0 K^+) + \sqrt{2}A(B^0 \rightarrow \pi^0 K^0) - A(B^+ \rightarrow \pi^+ K^0) = 0. \quad (1)$$

The four amplitudes may be written in terms of penguin (P_{tc} and P_{uc}), tree (T), color-suppressed (C), annihilation (A), color-favored electroweak penguin (P_{EW}), and color-suppressed electroweak penguin (P_{EW}^C) contributions [2]:

$$-A(B^0 \rightarrow \pi^- K^+) = P_{tc} + P_{uc} + T + \frac{2}{3}P_{EW}^C, \quad (2)$$

$$-\sqrt{2}A(B^+ \rightarrow \pi^0 K^+) = P_{tc} + P_{uc} + T + C + A + P_{EW} + \frac{2}{3}P_{EW}^C, \quad (3)$$

$$\sqrt{2}A(B^0 \rightarrow \pi^0 K^0) = P_{tc} + P_{uc} - C - P_{EW} - \frac{1}{3}P_{EW}^C, \quad (4)$$

$$A(B^+ \rightarrow \pi^+ K^0) = P_{tc} + P_{uc} + A - \frac{1}{3}P_{EW}^C. \quad (5)$$

The two amplitudes P_{tc} and P_{uc} behave like isoscalars ($\Delta I = 0$) while the remaining terms are mixtures of isoscalar and isovector ($\Delta I = 1$). The terms P_{tc} , P_{EW} and P_{EW}^C involve a CKM factor $V_{tb}^* V_{ts}$ with weak phase π , while P_{uc} , T , C and A contain $V_{ub}^* V_{us}$ with phase γ . Each of these amplitudes has its own unknown strong phase, but some strong phases can be related to each other approximately. Using flavor SU(3) symmetry, the amplitudes P_{EW} and P_{EW}^C are given to a good approximation in terms of T and C [3]:

$$P_{EW} = \frac{3}{4} \frac{c_9 + c_{10}}{c_1 + c_2} R(T + C) + \frac{3}{4} \frac{c_9 - c_{10}}{c_1 - c_2} R(T - C) \approx \frac{3}{2} \frac{c_9 + c_{10}}{c_1 + c_2} RT, \quad (6)$$

$$P_{EW}^C = \frac{3}{4} \frac{c_9 + c_{10}}{c_1 + c_2} R(T + C) - \frac{3}{4} \frac{c_9 - c_{10}}{c_1 - c_2} R(T - C) \approx \frac{3}{2} \frac{c_9 + c_{10}}{c_1 + c_2} RC. \quad (7)$$

Here, c_i are Wilson coefficients [4] obeying $(c_9 + c_{10})/(c_1 + c_2) \approx (c_9 - c_{10})/(c_1 - c_2)$, while $R \equiv |(V_{tb}^* V_{ts})/(V_{ub}^* V_{us})| = 48.9 \pm 1.6$ [5]. The proportionality coefficient relating P_{EW} to T and P_{EW}^C to C is numerically of order one and negative: $\delta_{EW} \equiv (3/2)[(c_9 + c_{10})/(c_1 + c_2)]R = -0.60 \pm 0.02$. SU(3) breaking introduces a theoretical error of about 10% in the magnitude of this coefficient and an error of 5° in its strong phase [6].

The decays $B \rightarrow \pi K$ provide nine measurements: four branching fractions $\mathcal{B}$, four direct CP asymmetries A_{CP} , and the mixing-induced CP asymmetry S_{CP} in $B^0 \rightarrow \pi^0 K^0$. Assuming values for the weak phases β and γ as determined in a global CKM fit [5], these observables can be expressed in terms of nine parameters: the magnitude of the five independent amplitudes, P_{tc} , P_{uc} , T , C , A , and their four

Table 1: Branching fractions and CP asymmetries for $B \rightarrow \pi K$ decays, as of today and for early 2007 (in parentheses) [13].

Mode	$\mathcal{B} (10^{-6})$	A_{CP}	S_{CP}
$B^0 \rightarrow \pi^- K^+$	19.4 ± 0.6 (19.7 ± 0.6)	$-0.098^{+0.012}_{-0.011}$ (-0.093 ± 0.015)	
$B^+ \rightarrow \pi^0 K^+$	12.9 ± 0.6 (12.8 ± 0.6)	0.050 ± 0.025 (0.047 ± 0.026)	
$B^0 \rightarrow \pi^0 K^0$	9.8 ± 0.6 (10.0 ± 0.6)	-0.01 ± 0.10 (-0.12 ± 0.11)	0.57 ± 0.17 (0.33 ± 0.21)
$B^+ \rightarrow \pi^+ K^0$	23.1 ± 1.0 (23.1 ± 1.0)	0.009 ± 0.025 (0.009 ± 0.025)	

relative strong phases. Neglecting the amplitude A , which vanishes to leading order in $1/m_b$ and α_s [7], one may perform a best fit with two degrees of freedom. Such a fit was made about two years ago [8] with data available in early 2007. The fit neglected P_{uc} but kept γ as a free parameter which was extracted successfully in agreement with its determination in the CKM fit. A good fit, corresponding to $\chi^2_{min}/d.o.f. = 1/3$ (80% c.l.), required $|C/T| = 1.6 \pm 0.3$. In contrast, values of $|C/T|$ calculated in QCD, using a heavy-quark expansion and various assumptions about Λ_{QCD}/m_b corrections, do not exceed 0.6 [9, 10]. The larger value of $|C/T|$ obtained in the fit [8] was thus considered a possible indication for New Physics (NP) [11]. (Another good fit to the data, involving a sizable contribution from P_{uc} , gave a lower value $|C/T| = 0.8 \pm 0.1$. However, its value of γ was in disagreement with the global CKM fit.)

Recently, an update of the fit to $B \rightarrow \pi K$ data was performed using the latest information as of early 2009 [12]. The current data are summarized in Table 1 [13], compared with the data of two years ago in parentheses. The only significant change occurred in the central values of the two CP asymmetries A_{CP} and S_{CP} in $B^0 \rightarrow \pi^0 K^0$, which involve the largest experimental errors. The current experimental situation with respect to the need for NP has now become less clear. Although the best fit [12], including P_{uc} as a parameter and using γ as an input, is relatively poor [corresponding to $\chi^2_{min}/d.o.f. = 3.2/2$ (20%)], it now requires $|C/T| = 0.58 \pm 0.24$. This value is consistent with QCD calculations (within their uncertainties) and with flavor SU(3) fits combining $B \rightarrow \pi K$ and $B \rightarrow \pi\pi$ data [14].

A test for the CKM framework based on the asymmetry $S_{CP}(\pi^0 K^0)$ was suggested in Ref. [15]. Assuming flavor SU(3) in order to obtain the quantity $|T + C|$ from the decay rate for $B^+ \rightarrow \pi^+ \pi^0$ [16], the asymmetry $S_{CP}(\pi^0 K^0)$ can be determined using as inputs this rate measurement and all the $B \rightarrow \pi K$ measurements except $S_{CP}(\pi^0 K^0)$. It was noted, however, that this test is sensitive to input values of $\mathcal{B}(B^0 \rightarrow \pi^0 K^0)$, $\mathcal{B}(B^0 \rightarrow \pi^- K^+)$ and to SU(3) breaking in δ_{EW} [17].

The branching fractions in Table 1 and the lifetime ratio [13] $\tau_+/\tau_0 \equiv \tau(B^+)/\tau(B^0) = 1.073 \pm 0.008$ imply that the corresponding decay widths are approximately in the ratio 2:1:1:2. This indicates the dominance of the isospin-preserving ($\Delta I = 0$) amplitude P_{tc} . Indeed, in the best fit of Ref. [12], $|T|$ is about 12% of $|P_{tc}|$ while other contributions are smaller. Assuming that C is suppressed relative to T and can be neglected, it was shown [18] that the rate differences

$$\Delta(\pi^- K^+) \equiv \Gamma(\bar{B}^0 \rightarrow \pi^+ K^-) - \Gamma(B^0 \rightarrow \pi^- K^+) \quad (8)$$

and

$$2\Delta(\pi^0 K^+) \equiv 2[\Gamma(B^- \rightarrow \pi^0 K^-) - \Gamma(B^+ \rightarrow \pi^0 K^+)] \quad (9)$$

should be approximately equal. As the CP-averaged rate for $B^0 \rightarrow \pi^- K^+$ is about double that for $B^+ \rightarrow \pi^0 K^+$, this translates to a prediction of approximately equal CP asymmetries,

$$A_{CP}(\pi^- K^+) \approx A_{CP}(\pi^0 K^+) . \quad (10)$$

The current measured asymmetries given in Table 1 differ by more than 5σ . This has occasionally been taken to indicate the presence of NP in $B \rightarrow \pi K$ [19]. It was noted, however [20], that a nonzero contribution of C at a level suggested in SU(3) fits [14] (and recently in the $B \rightarrow \pi K$ fit [12]) could account for this difference. A necessary requirement is that the strong phase difference $\arg(C/T)$ is large and negative [21]. A large negative phase has been obtained in flavor SU(3) fits to $B \rightarrow \pi K$ and $B \rightarrow \pi\pi$ data [14]. A phase of about -130° , obtained in the best fit in Ref. [12], and the central value $|C/T| = 0.58$ account well for the difference between the two asymmetries. While this magnitude of C/T can be accounted for in QCD calculations, its large negative phase seems a problem for certain QCD calculations [7, 9] but not for others [10].

A very robust sum rule was suggested a few years ago [22] combining all four CP rate asymmetries in $B \rightarrow \pi K$:

$$\Delta(\pi^- K^+) + \Delta(\pi^+ K^0) \approx 2\Delta(\pi^0 K^+) + 2\Delta(\pi^0 K^0) , \quad (11)$$

where $\Delta(\pi^+ K^0)$ and $\Delta(\pi^0 K^0)$ are defined similarly to $\Delta(\pi^- K^+)$ and $\Delta(\pi^0 K^+)$ in Eqs. (8) and (9). This approximate sum rule is based on the amplitude isospin relation [Eq. (1)] and its CP conjugate, in which the isoscalar ($\Delta I = 0$) and isovector ($\Delta I = 1$) parts vanish separately. Consider the difference between the left-hand side of Eq. (11) and its right-hand side. The leading terms in this difference involve interference between the dominant isoscalar amplitude P_{tc} and a linear combination of smaller isoscalar and isovector amplitudes, each of which vanishes because of Eq. (1). The remaining terms arise from the interference of smaller $\Delta I = 1$ amplitudes with one another. These terms were found to vanish in the flavor SU(3) and heavy-quark limits [22]. A simplified and slightly less precise version of the sum rule [Eq. (11)], which assumes that the rates in Table 1 are in the ratio 2:1:1:2, relates the CP asymmetries directly:

$$A_{CP}(\pi^- K^+) + A_{CP}(\pi^+ K^0) \simeq A_{CP}(\pi^0 K^+) + A_{CP}(\pi^0 K^0) . \quad (12)$$

The sum rule in Eq. (11) holds also in the presence of a $\Delta I = 0$ NP amplitude which can be absorbed in P_{tc} in the above argument. Furthermore, since the sum rule can only be violated by terms which are quadratic in $\Delta I = 1$ amplitudes, a substantial violation of the sum rule requires isovector NP amplitudes which are not much smaller than P_{tc} . This will be the basis of our following diagnostic for NP in $B \rightarrow \pi K$ decays.

While any evidence for NP in $B \rightarrow \pi K$ is currently weak, suppose that NP is present in these decays. It was argued in Ref. [23] that all NP strong phases are negligible. Given that all strong phases are equal, there are two classes of NP amplitudes contributing to $B \rightarrow \pi K$, differing only in their color structure [24]:

$$\mathcal{A}^q e^{i\Phi_q} \equiv \sum \langle \pi K | \bar{s}_\alpha \Gamma_i b_\alpha \bar{q}_\beta \Gamma_j q_\beta | B \rangle , \quad (13)$$

$$\mathcal{A}^{C,q} e^{i\Phi_q^C} \equiv \sum \langle \pi K | \bar{s}_\alpha \Gamma_i b_\beta \bar{q}_\beta \Gamma_j q_\alpha | B \rangle , \quad (14)$$

where $\Gamma_{i,j}$ represent Lorentz structures and $q = u, d$. (Despite the index C standing for color-suppression, the matrix elements $\mathcal{A}^{C,q} e^{i\Phi_q^C}$ are not necessarily smaller than $\mathcal{A}^q e^{i\Phi_q}$.) Here, Φ_q and Φ_q^C are the NP weak phases; the strong phases are taken to be zero. The NP amplitudes $\mathcal{A}^q e^{i\Phi_q}$ and $\mathcal{A}^{C,q} e^{i\Phi_q^C}$ are equivalent to the amplitudes ΔP_q and ΔP_q^c defined in Ref. [25].

There are three NP matrix elements which contribute to the $B \rightarrow \pi K$ amplitudes: $\mathcal{A}^{comb} e^{i\Phi} \equiv -\mathcal{A}^u e^{i\Phi_u} + \mathcal{A}^d e^{i\Phi_d}$, $\mathcal{A}^{C,u} e^{i\Phi_u^C}$, and $\mathcal{A}^{C,d} e^{i\Phi_d^C}$ [24]. (These are equivalent to the three independent NP combinations, $-\Delta P_u + \Delta P_d$, ΔP_u^c , and ΔP_d^c , contributing to $B \rightarrow \pi K$ amplitudes as discussed in Ref. [25].) The first operator corresponds to including NP only in the color-favored electroweak penguin amplitude: $\mathcal{A}^{comb} e^{i\Phi} \equiv -P_{EW,NP} e^{i\Phi_{EW}}$. Nonzero values of $\mathcal{A}^{C,u} e^{i\Phi_u^C}$ and/or $\mathcal{A}^{C,d} e^{i\Phi_d^C}$ imply the inclusion of NP in both the gluonic and color-suppressed electroweak penguin amplitudes, $P_{NP} e^{i\Phi_P}$ and $P_{EW,NP}^C e^{i\Phi_{EW}^C}$, respectively [26]:

$$P_{NP} e^{i\Phi_P} \equiv \frac{1}{3} \mathcal{A}^{C,u} e^{i\Phi_u^C} + \frac{2}{3} \mathcal{A}^{C,d} e^{i\Phi_d^C} , \quad (15)$$

$$P_{EW,NP}^C e^{i\Phi_{EW}^C} \equiv \mathcal{A}^{C,u} e^{i\Phi_u^C} - \mathcal{A}^{C,d} e^{i\Phi_d^C} . \quad (16)$$

Thus, NP in $B \rightarrow \pi K$ is of one of the above three varieties. It can affect the gluonic penguin amplitude ($P_{NP} e^{i\Phi_P}$), the color-favored electroweak penguin amplitude ($P_{EW,NP} e^{i\Phi_{EW}}$), or the color-suppressed electroweak penguin amplitude ($P_{EW,NP}^C e^{i\Phi_{EW}^C}$). These three NP amplitudes are added to P_{tc} , P_{EW} and P_{EW}^C , respectively. In the first case, the NP is $\Delta I = 0$ and, as mentioned, will not affect the sum rule of Eq. (11). In the other two cases, the NP breaks isospin by one unit, so that the sum rule will be violated. Ref. [12] performed fits with each of the three NP operators. In the SM-like fit it was found that it is not possible to constrain NP in the $\Delta I = 0$ gluonic penguin ($P_{NP} e^{i\Phi_P}$). This is not surprising – it is the same in $B \rightarrow \pi\pi$ decays [27]. However, the two non-SM-like fits did indeed involve substantial contributions from NP: $|P_{EW,NP}/P_{tc}| = 0.4$, $|P_{EW,NP}^C/P_{tc}| = 0.3$. They had lower values of $\chi^2/d.o.f.$ (0.4/2 and 2.5/2) than the SM-like fit (3.6/2). One potentially troublesome point is that they had very large values of $|C/T|$ (6 ± 11

and 4.9 ± 3.8). However, the errors are also quite large, so this might not be a real problem. The key point here is that these two NP operators are of the type $\Delta I = 1$, and so can violate the sum rules of Eqs. (11) and (12).

Using the CP asymmetries [all except for $A_{CP}(B^0 \rightarrow \pi^0 K^0)$] and branching ratios in Table 1, and translating ratios of branching ratios into ratios of rates using the lifetime ratio $\tau_+/\tau_0 = 1.073 \pm 0.008$ [13], the sum rule of Eq. (11) predicts

$$A_{CP}(B^0 \rightarrow \pi^0 K^0) = -0.149 \pm 0.044 , \quad (17)$$

more than 3σ away from zero. The $B^0 \rightarrow \pi^0 K^0$ mode is thus predicted to exhibit the largest CP asymmetry of any of the four $B \rightarrow \pi K$ modes.

If one uses the simplified version of the sum rule [Eq. (12)], which assumes the rates in Table 1 are in the ratio 2:1:1:2, the CP asymmetries are related directly:

$$A_{CP}(\pi^0 K^0) = A_{CP}(\pi^- K^+) + A_{CP}(\pi^+ K^0) - A_{CP}(\pi^0 K^+) = -0.139 \pm 0.037 . \quad (18)$$

Indeed, the SM-like fit in Ref. [12] finds $A_{CP}(\pi^0 K^0) = -0.12$. (We restrict our attention to those fits which include the constraints on CKM phases [13] $\beta = (21.66^{+0.95}_{-0.85})^\circ$ and $\gamma = (66.8^{+5.4}_{-3.8})^\circ$.)

On the other hand, in the other two fits of Ref. [12], $\Delta I = 1$ NP amplitudes are involved, which are much more significant than those in the SM, and very large values of $|C|$ are obtained. This results in a crucial difference with the SM fit: both of these fits predict $A_{CP}(\pi^0 K^0) = -0.03$. We thus see that it is possible to detect the presence of NP with a precise measurement of $A_{CP}(\pi^0 K^0)$.

Another sum rule involving decay rates instead of CP asymmetries is, in principle, sensitive to NP in $\Delta I = 1$ amplitudes [18, 30]. In terms of branching ratios $\mathcal{B}(B \rightarrow f) \equiv \mathcal{B}(f)$ it is expressed as

$$2\mathcal{B}(\pi^0 K^+) + 2(\tau_+/\tau_0)\mathcal{B}(\pi^0 K^0) = (\tau_+/\tau_0)\mathcal{B}(\pi^- K^+) + \mathcal{B}(\pi^+ K^0) . \quad (19)$$

The difference between the left- and right-hand sides is quadratic in $\Delta I = 1$ amplitudes, expected to be small in the Standard Model. A discussion of the terms violating this sum rule can be found in Ref. [25], which also discusses NP in $B \rightarrow \pi K$ decays.

The experimental branching ratios in Table 1 satisfy the sum rule at the 1.4σ level, with the (left, right)-hand sides giving $(46.8 \pm 1.8, 43.9 \pm 1.2)$ in units of 10^{-6} . The three fits of Ref. [12] satisfy the sum rule as well or better, as shown in Table 2. A large $P_{EW,NP}$ amplitude in Fit 2 is nearly cancelled by a large C contribution in a delicate way so as to preserve the sum rule.

One may ask if the large value of $|C/T|$ in Fits 2 and 3 is obligatory. Performing the same fits but constraining $|C/T| = 0.5$, one finds results summarized in Table 2. The number of degrees of freedom with the constraint $|C/T| = 0.5$ is 3, while without the constraint it is 2. It appears that a large value of $|C|$, along with a sizable NP amplitude, is responsible for moving the predicted $A_{CP}(\pi^0 K^0)$ away from its Standard Model value of -0.15 .

The difference between the violation of the rate sum rule [Eq. (19)] and the asymmetry sum rule [Eq. (11)] in the presence of NP can be seen as follows. Define

Table 2: Comparison of fits of Ref. [12] with and without constraint $|C/T| = 0.5$. Prediction for (a) $A_{CP}(\pi^0 K^0)$, (b) Eq. (19), l.h.s; (c) Eq. (19), r.h.s.

Fit	With $ C/T = 0.5$				Without $ C/T = 0.5$			
	$\chi^2/d.o.f.$	(a)	(b)	(c)	$\chi^2/d.o.f.$	(a)	(b)	(c)
1	3.7/3	-0.11	44.9	44.8	3.6/2	-0.12	44.9	44.8
2	3.0/3	-0.12	47.1	43.2	0.4/2	-0.03	47.0	43.8
3	3.8/3	-0.12	44.9	44.8	2.5/2	-0.03	45.3	44.6

$P_1 \equiv P_{EW,NP}$ and $P_2 \equiv P_{EW,NP}^C$ and assume they are large, neglecting the violation of the two sum rules by small SM contributions (which is a good approximation).

The terms violating the rate sum rule are

$$2[|P_1|^2 + |P_1||P_2|\cos(\delta_1 - \delta_2)\cos(\phi_1 - \phi_2)] , \quad (20)$$

where δ_1, δ_2 and ϕ_1, ϕ_2 are the strong and weak phases of P_1 and P_2 . This leads to a violation of the rate sum rule [Eq. (19)] in the case $P_1 \neq 0, P_2 = 0$ (Fit 2 of Ref. [12]) but to no violation in the case $P_1 = 0, P_2 \neq 0$ (Fit 3 of Ref. [12]).

On the other hand, the term violating the asymmetry sum rule is

$$2|P_1||P_2|\sin(\delta_1 - \delta_2)\sin(\phi_1 - \phi_2) . \quad (21)$$

In Ref. [12] it was assumed that $\delta_1 = \delta_2$, so the asymmetry sum rule was not violated by this term. Also, the cases $P_1 = 0, P_2 \neq 0$ and $P_1 \neq 0, P_2 = 0$ were considered separately. In these cases the sum rule would hold even when $\delta_1 \neq \delta_2$.

The reason for the violation of the asymmetry sum rule in the two fits (Fit 2 and Fit 3) of Ref. [12] without a constraint on $|C/T|$ (seventh column, second and third rows of Table 2), giving $A_{CP} = -0.03$ instead of -0.12 , is that C was also large in both fits, so the sum rule was violated by contributions

$$4|P_1||C|\sin(\delta_1 - \delta_C)\sin(\phi_1 - \gamma) \quad (\text{Fit 2}) , \quad (22)$$

$$2|P_2||C|\sin(\delta_2 - \delta_C)\sin(\phi_2 - \gamma) \quad (\text{Fit 3}) . \quad (23)$$

To sum up, the asymmetry sum rule [Eq. (11)] can be violated significantly (when assuming $|C/T|$ is not very large) only by taking $P_1 \neq 0, P_2 \neq 0, \delta_1 \neq \delta_2$, and $\phi_1 \neq \phi_2$. In other words, violation of the sum rule requires both $P_{EW,NP}$ and $P_{EW,NP}^C$ to be present and both strong- and weak-phase differences of these amplitudes to be non-negligible. One may imagine a situation in which these circumstances hold leading to an asymmetry $A_{CP} = -0.03$ in $B^0 \rightarrow \pi^0 K^0$ as discussed above.

Although we recognize the difficulty of the measurement, we thus urge a reduction of the present experimental error on $A_{CP}(\pi^0 K^0)$, whose world average is -0.01 ± 0.10 . The BaBar and Belle contributions to this average are compared in Table 3. The error must be reduced by at least a factor of three in order to

Table 3: Comparison of BaBar [28] and Belle [29] values of $A_{CP}(B^0 \rightarrow \pi^0 K^0)$.

Source	$N(B\bar{B})$ (M)	A_{CP}
BaBar [28]	467	$-0.13 \pm 0.13 \pm 0.03$
Belle [29]	657	$0.14 \pm 0.13 \pm 0.06$
Average [13]	1124	-0.01 ± 0.10

determine whether new physics is generating a large $\Delta I = 1$ transition amplitude. An experiment collecting of the order of 10^{10} $B\bar{B}$ pairs ought to be able to make the necessary distinction.

Acknowledgments: This work was financially supported in part by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) No. KRF-2007-359-C00009 (SB), by the National Science Council of Taiwan, R.O.C. under Grant No. NSC 97-2112-M-008-002-MY3 (CC), by NSERC of Canada (DL), and by the United States Department of Energy through Grant No. DE FG02 90ER40560 (JR).

References

- [1] Y. Nir and H. R. Quinn, Phys. Rev. Lett. **67** (1991) 541; M. Gronau, Phys. Lett. B **265** (1991) 389.
- [2] M. Gronau, O. F. Hernandez, D. London and J. L. Rosner, Phys. Rev. D **50** (1994) 4529 [arXiv:hep-ph/9404283], Phys. Rev. D **52** (1995) 6374 [arXiv:hep-ph/9504327].
- [3] M. Neubert and J. L. Rosner, Phys. Lett. B **441** (1998) 403 [arXiv:hep-ph/9808493], Phys. Rev. Lett. **81** (1998) 5076 [arXiv:hep-ph/9809311]; M. Gronau, D. Pirjol and T. M. Yan, Phys. Rev. D **60** (1999) 034021 [Erratum-ibid. D **69** (2004) 119901] [arXiv:hep-ph/9810482]; M. Imbeault, A. L. Lemerle, V. Page and D. London, Phys. Rev. Lett. **92** (2004) 081801 [arXiv:hep-ph/0309061].
- [4] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. **68** (1996) 1125 [arXiv:hep-ph/9512380].
- [5] J. Charles *et al.* [CKMfitter Group], Eur. Phys. J. C **41** (2005) 1 [arXiv:hep-ph/0406184]. Updated results are available at <http://ckmfitter.in2p3.fr>.
- [6] M. Neubert, JHEP **9902** (1999) 014 [arXiv:hep-ph/9812396].

- [7] C. W. Bauer, D. Pirjol, I. Z. Rothstein and I. W. Stewart, Phys. Rev. D **70** (2004) 054015 [arXiv:hep-ph/0401188]; C. W. Bauer, I. Z. Rothstein and I. W. Stewart, Phys. Rev. Lett. **94** (2005) 231802 [arXiv:hep-ph/0412120].
- [8] S. Baek and D. London, Phys. Lett. B **653** (2007) 249 [arXiv:hep-ph/0701181].
- [9] M. Beneke and S. Jager, Nucl. Phys. B **751** (2006) 160 [arXiv:hep-ph/0512351].
- [10] H. n. Li and S. Mishima, arXiv:0901.1272 [hep-ph].
- [11] See also A. J. Buras, R. Fleischer, S. Recksiegel and F. Schwab, Phys. Rev. Lett. **92** (2004) 101804 [arXiv:hep-ph/0312259]; Nucl. Phys. B **697** (2004) 133 [arXiv:hep-ph/0402112].
- [12] S. Baek, C. W. Chiang and D. London, arXiv:0903.3086 [hep-ph], to be published in Physics Letters B.
- [13] Updated results and references are tabulated periodically by the Heavy Flavor Averaging Group: <http://www.slac.stanford.edu/xorg/hfag/rare>. See, e.g., E. Barberio *et al.* arXiv: 0704.3575v1, <http://www.slac.stanford.edu/xorg/hfag/results/>.
- [14] C. W. Chiang, M. Gronau, J. L. Rosner and D. A. Suprun, Phys. Rev. D **70** (2004) 034020 [arXiv:hep-ph/0404073]; C. W. Chiang and Y. F. Zhou, JHEP **0612** (2006) 027 [arXiv:hep-ph/0609128].
- [15] R. Fleischer, S. Jager, D. Pirjol and J. Zupan, Phys. Rev. D **78** (2008) 111501 [arXiv:0806.2900 [hep-ph]].
- [16] M. Gronau, J. L. Rosner and D. London, Phys. Rev. Lett. **73** (1994) 21 [arXiv:hep-ph/9404282].
- [17] M. Gronau and J. L. Rosner, Phys. Lett. B **666** (2008) 467 [arXiv:0807.3080 [hep-ph]].
- [18] M. Gronau and J. L. Rosner, Phys. Rev. D **59** (1999) 113002 [arXiv:hep-ph/9809384].
- [19] See, for instance, S.-W. Lin et al. (Belle collaboration), Nature **452** (2008) 332; M. Peskin, *ibid.* **452** (2008) 293.
- [20] M. Gronau and J. L. Rosner, Phys. Rev. D **74** (2006) 057503 [arXiv:hep-ph/0608040].
- [21] M. Gronau and J. L. Rosner, Phys. Lett. B **644** (2007) 237 [arXiv:hep-ph/0610227].
- [22] M. Gronau, Phys. Lett. B **627** (2005) 82 [arXiv:hep-ph/0508047].

- [23] A. Datta and D. London, Phys. Lett. B **595** (2004) 453 [arXiv:hep-ph/0404130].
- [24] A. Datta, M. Imbeault, D. London, V. Page, N. Sinha and R. Sinha, Phys. Rev. D **71** (2005) 096002 [arXiv:hep-ph/0406192].
- [25] M. Gronau and J. L. Rosner, Phys. Rev. D **71** (2005) 074019 [arXiv:hep-ph/0503131].
- [26] S. Baek, JHEP **0607** (2006) 025 [arXiv:hep-ph/0605094].
- [27] S. Baek, F. J. Botella, D. London and J. P. Silva, Phys. Rev. D **72** (2005) 036004 [arXiv:hep-ph/0506075].
- [28] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **79** (2009) 052003 [arXiv:0809.1174 [hep-ex]].
- [29] I. Adachi *et al.* [Belle Collaboration], arXiv:0809.4366 [hep-ex].
- [30] H. J. Lipkin, Phys. Lett. B **445** (1999) 403 [arXiv:hep-ph/9810351].